

AGELESS IMAGE WELLNESS AND BEAUTY

AFTERCARE INSTRUCTIONS FOR KYBELLA TREATMENT:



Apply ice to the treatment area for up to twenty minutes to decrease discomfort.



Bruising can take up to two weeks to resolve. After the first thirty six hours you can use concealer or make-up to decrease their appearance. You can also apply Arnica gel to the bruised areas to decrease healing time.



DO NOT TAKE ANTI-INFLAMMATORY MEDICATIONS. These include Motrin, or Ibuprofen, Aspirin and Aleve for at least 48 hours.



There will be a large amount of swelling for the first twenty four to forty eight hours following treatment. This swelling can be significant and feel jelly or liquid like.



The swelling can give you a “wobbly” sensation, in the treated area, for five to seven days.



The jelly like feel can change to a hard lump within the treated area. This happens in the first week and is a normal response. It is a sign your results will be GREAT.



The swelling associated with treatment will resolve slowly over the next four to eight weeks.



EXPECT NUMBNESS IN THE TREATED AREA. This can be profound and last up to three months. It will usually resolve in four to six weeks but can last longer.



If not done at the time of treatment please call the office and schedule your two week follow-up appointment. This is complimentary, we want to make sure you are satisfied with your care and you are progressing appropriately



Treatment can cause temporary asymmetry of your smile. If this occurs call the office for an appointment to be evaluated.



THANK YOU FOR CHOOSING AGELESS IMAGE TO CARE FOR YOU TODAY! WE APPRECIATE YOU AND LOOK FORWARD TO HELPING YOU ACHIEVE YOUR GOALS.

PURSUING BEAUTY THAT IS MORE THAN SKIN DEEP